

# Motor and cognitive functions of the ventral premotor cortex

## Giacomo Rizzolatti\*<sup>‡</sup>, Leonardo Fogassi<sup>†</sup> and Vittorio Gallese\*

Recent data show that the ventral premotor cortex in both humans and monkeys has motor and cognitive functions. The cognitive functions include space perception, action understanding and imitation. The data also show a clear functional homology between monkey area F5 and human area 44. Preliminary evidence suggests that the ventral part of the lateral premotor cortex in humans may correspond to monkey area F4. A tentative map of the human lateral premotor areas founded on the reviewed evidence is presented.

### Addresses

\*Istituto di Fisiologia Umana and <sup>†</sup>Dipartimento di Psicologia, Università di Parma, Via Volturno 39, I-43100 Parma, Italy  
<sup>‡</sup>e-mail: fisioium@symbolic.pr.it

**Current Opinion in Neurobiology** 2002, **12**:149–154

0959-4388/02/\$ – see front matter  
 © 2002 Elsevier Science Ltd. All rights reserved.

**Published online 4th March 2002**

### Abbreviations

<b>AIP</b>	anterior intraparietal area
<b>dPM</b>	dorsal premotor area
<b>FEF</b>	frontal eye field
<b>fMRI</b>	functional magnetic resonance imaging
<b>RFs</b>	receptive fields
<b>SII</b>	secondary somatosensory cortex
<b>VIP</b>	ventral intraparietal area
<b>vPM</b>	ventral premotor area

### Introduction

The premotor cortex of the monkey is typically subdivided into two large sectors: the ventral (vPM) and the dorsal (dPM) premotor cortex. In turn, each of these sectors is formed by two areas, one located rostrally and the other caudally (Figure 1a). No distinction between dorsal and ventral premotor cortex has classically been made in humans, the cortex lying on the cortical lateral surface being dubbed lateral area 6. A distinction, however, between rostral and caudal premotor sectors is apparent in humans. The dorsal part of lateral area 6 consists of two areas: 6a $\alpha$  and 6a $\beta$  [1]. They are indicated in light pink and blue, respectively, in Figure 1b. Two areas similarly form the ventral sector of lateral area 6: 6a $\alpha$  and area 44. They are indicated in red and dark pink, respectively, in Figure 1b. The homology between area 44 in humans and the rostral part of vPM (or F5) in monkeys was noted by Von Bonin and Bailey [2] and recently confirmed by Petrides and Pandya [3].

Here, we review recent monkey and human studies that address the functional role of ventral sectors of premotor cortex (area 44 included) in action organization, motor imagery, and action understanding. Studies on language are not reviewed.

### Caudal ventral premotor cortex in monkeys and humans

#### Caudal ventral premotor cortex in monkeys: F4

Area F4 is the frontal node of a cortical circuit that includes the ventral intraparietal area (VIP), the intraparietal sector of area PE (PEip), and the secondary somatosensory cortex (SII) [4<sup>•</sup>]. This circuit transforms specific positions in the space around the monkey (peripersonal space) into arm, neck, and face/mouth movements. This circuit is also involved in space perception [5].


F4 neurons discharge according to specific body part movements. In addition, most of these neurons respond to somatosensory stimuli. Some also respond to visual and auditory stimuli [6–9]. F4 visual receptive fields (RFs) are located around the tactile RF and, typically, extend in depth up to ~30 cm from it [6–8]. Their position is independent of eye position and is anchored to a particular body part [10]. When the body part is moved, the RF moves with it [8]. Recent data showed that F4 contains representations of wrist movements directed to specific space positions [11<sup>•</sup>]. Interestingly, most F4 neurons that respond to wrist movements code actions in extrinsic (spatial) coordinates, rather than in intrinsic coordinates, as previously described for F1 [12]. This study [11<sup>•</sup>] elegantly confirmed the notion that area F4 codes goal-directed actions, the goal being represented by spatial locations.

#### Lesion studies

Ablation of F4 (including portions of F5) produces hemispatial neglect in the monkey, as well as reaching deficits. The neglect is circumscribed to peripersonal space [13]. Selective neglect for peripersonal space may occur also in humans [14]. A recent study by Berti and Frassinetti [15<sup>•</sup>] confirmed this finding. They described a patient who, after a right hemisphere stroke, exhibited selective peripersonal neglect. The near/far dissociation was clear in line bisection tasks that were impaired in the near space, but virtually normal in the far space. This was observed when the patient bisected lines using a light projection pen. However, when the patient acted on far stimuli using a stick, neglect reappeared. Berti and Frassinetti [15<sup>•</sup>] explained this result by referring to neurophysiological data obtained by Iriki *et al.* [16<sup>••</sup>], who showed that the use of a tool extends the monkey peripersonal space. In the case of the patient, the use of a stick extended his peripersonal space so as to include all the space between the body and the stimulus. As a consequence, far space was remapped as near space and the neglect reappeared. Maravita *et al.* [17] and Farnè and Ladavas [18] reported similar findings.

Prompted by the observation that, in the monkey, F4 visual RFs move with the body part to which they are

Figure 1



Ventral premotor cortex in monkeys and humans. (a) Lateral view of the monkey brain showing the cytoarchitectonic parcellation of the motor cortex and the posterior parietal cortex. Matching colors indicate each motor area and its corresponding parietal input (area VIP, buried inside IP, is not shown). The motor area F7, receiving its main input from the prefrontal lobe, is indicated in blue. Areas F2 and F7 are often referred to as dPM; areas F4 and F5 form the vPM. AI, inferior arcuate sulcus; AS, superior arcuate sulcus; C, central sulcus; DLPF<sub>d</sub>, dorsolateral prefrontal cortex dorsal part; DLPF<sub>v</sub>, dorsolateral prefrontal cortex ventral part; IP, intraparietal sulcus; L, lateral fissure; Lu, lunate sulcus; P, principal sulcus; SI, primary somatosensory cortex; ST, superior temporal sulcus. Modified with permission from [4\*]. (b) Lateral view of the human brain showing the cytoarchitectonic parcellation of Brodmann and the basal lines (thick lines) depicted by Talairach and Tournoux [50]. Thin horizontal lines indicate z values. Colored regions indicate: dPM 6 $\alpha$ , light pink; 6 $\beta$  (predPM [51\*]), blue; vPM (ventral area 6), red; area 44, dark pink; FEF, green. Modified with permission from [50].

Current Opinion in Neurobiology

anchored, a series of studies investigated cross-modal extinction in patients with mild neglect [19,20]. These experiments showed that the presentation of a visual stimulus near the patients' ipsilesional (normal) arm or hemiface caused an extinction of a tactile stimulus applied on the patients' contralesional (neglected) hand. However, when the visual stimulus was presented in the identical spatial location, but the body part was moved to another position, the extinguishing effect of vision was not observed or was markedly reduced. Peripersonal cross-modal extinction was also recently observed when the ipsilesional visual stimulus was presented near the patient's hand seen in a mirror [21\*].

The lesions causing neglect in humans are always very large. Thus, although data supporting the existence of separate

systems for peripersonal and extrapersonal space are very robust, the localization of the human homologue of area F4 is difficult to determine on the basis of lesion data alone.

#### Caudal ventral premotor cortex in humans

Brain imaging experiments, in which the anatomical localization of arm movements was studied, were unsuccessful in localizing F4 in humans. The results showed only a strong activation of the primary motor cortex extending to the dorsal part of area 6 (stereotaxic coordinates  $z=+52$  and above, see [22–25]). Note, however, that these studies used rather simple motor tasks not requiring the spatial transformations that characterize the activity of F4 neurons in monkeys. Furthermore, head and combined head/mouth movements that should also be represented in the human

homologue of F4 were never tested, due to the technical limitations of brain imaging techniques.

However, experiments using sensory stimulation suggest that an area homologous to monkey F4 exists in humans. Particularly interesting in this respect is a recent functional magnetic resonance imaging (fMRI) study [26\*\*]. Bremner *et al.* [26\*\*] attempted to localize the multimodal tactile, auditory and visual cortical areas in humans. Tactile stimuli (applied on the forehead), moving visual stimuli, and auditory stimuli producing the illusion of sound movement were presented to human subjects. Multimodal convergence was found in three sites: the depth of the intraparietal sulcus, SII, and the vPM at the border of ventral areas 6 and 44. The authors propose that the active vPM site probably corresponds to the monkey F4 sector where head movements are represented. Activation of ventral area 6 and area 44 was also recently found in a task in which participants had to judge the position of visual and somatosensory stimuli relative to their body midline [27]. The authors of this study suggested that these two areas intervene in cognitive spatial tasks.

Another experiment that supports the existence of a human homologue of F4 is that of Buccino *et al.* [28\*\*]. Here, participants were asked to observe goal-related foot, hand and mouth actions. The observation of arm reaching to grasp actions led to the activation of two premotor foci, one located in area 44 (see below), and one located ventral ( $x=-56$ ;  $y=-4$ ;  $z=+44$ ) to the activations observed during the execution of intransitive arm movements.

Summing up, it appears that humans may possess an area homologous to monkey area F4 (see Figure 1b). In this area, arm reaching movements are located dorsally [28\*\*], whereas head movements are located ventrally [26\*\*]. Note that this proposed homology is in accord with the location of the frontal eye field (FEF) in the two species. In monkeys, FEF is located immediately rostral to area F4 (see Figure 1a). In humans, the z coordinates of FEF range between +44 and +51 [29]. Our suggestion is that the upper limit of FEF in humans marks the border between vPM and dPM (see Figure 1b).

### Rostral ventral premotor cortex in monkeys and humans

Area F5 forms the rostral part of the monkey vPM. It consists of two main sectors: one, F5c, located on the dorsal convexity, the other, F5ab, on the posterior bank of the inferior arcuate sulcus. Both sectors receive afferents from SII and area PF of the parietal lobe. In addition, F5ab is the selective target of the anterior intraparietal area (AIP). Area F5 contains representations of hand and mouth movements. The two representations overlap to a considerable extent [4\*,5].

#### Canonical and mirror neurons

Recordings from single neurons in a natural context were particularly important for our understanding of the function

of F5. These studies showed that most F5 neurons code specific actions, rather than the single movements that form them. F5 neurons were thus subdivided into several action classes, for example, 'grasping', 'holding' or 'tearing' neurons [30]. Many F5 neurons respond to the presentation of visual stimuli (visuomotor neurons). Some of these respond to the presentation of graspable objects and were named 'canonical' F5 neurons (see [4\*] for review). Others discharge when the monkey observes another individual making an action in front of it. These neurons were named F5 'mirror' neurons [31,32]. Canonical neurons are mostly located in F5ab, whereas mirror neurons are found essentially in F5c.

On the basis of the functional properties of canonical F5 neurons, F5 was proposed to play a crucial role in transforming the visual properties of three-dimensional objects into hand shapes appropriate to interact with them. Recent inactivation experiments directly tested this proposal [33\*]. Monkeys were trained to reach for and grasp geometric solids of different size, shape and orientation. In separate sessions, F5ab, F5c and the hand field of F1 were reversibly inactivated. The results showed that after inactivation of F5ab (the sector where canonical neurons are located), the hand shaping that relies on the visual properties of the objects was markedly impaired. The monkeys were able to grasp the objects, but only after corrections made under tactile control.

It has been suggested that mirror neurons might be involved in the understanding of actions made by others [31,32]. Usually an action is recognized even when its final part is out of vision. A recent study investigated whether mirror neurons could form the basis of this capacity [34\*\*]. The experiment consisted of two conditions. In one, the monkey was shown a fully visible action directed towards an object. In the other, the same action was presented, but with its final, critical part (hand-object interaction) hidden behind a screen. The results showed that the majority of mirror neurons responded also in the hidden condition. In control experiments, the monkey saw the beginning of the same actions, but also knew that nothing was behind the screen. No response was obtained in this condition. These results clearly support the notion that mirror neurons mediate action understanding.

#### F5 and area 44 respond to hand actions

As mentioned above, cytoarchitecturally, the most likely human homologue of F5 is area 44 [2,3]. Functionally, however, whereas F5 has both a representation of hand and mouth actions [30], area 44 has been classically considered to be a speech area. Recent brain imaging experiments, by confirming previous indications (e.g. [35]), demonstrated beyond any doubt that area 44 is also related to hand actions [36,37\*,38,39\*,40,41\*].

Binkofski *et al.* [36] asked volunteers to manipulate multifaceted complex objects in one condition and a smooth

sphere in another. The complex manipulation task activated area 44 (−52, +8, +28), an area inside the intraparietal sulcus, and SII. This cortical network closely corresponds to the cortical circuit (AIP, SII, F5) for manipulation in the monkey [4\*]. Further support for the functional homology between F5 and area 44 comes from the study of Ehrsson *et al.* [37\*], in which they compared the cortical areas involved in different types of grips. They found that the focus of representation of precision grip is located in area 44 (−56, +12, +32). This finding fits with monkey data showing that precision grip is largely represented in F5, whereas only few F5 neurons discharge during whole hand prehension [30].

Area 44 motor representation is not only activated by actual movement execution, but also by imaging hand actions [42,43] or by hand mental rotation [44]. Gerardin *et al.* [39\*] showed that both transitive hand actions and the imagination of intransitive finger movements activate area 44 (−54, +12, +12). The activation was stronger during imagery than during actual execution. The prolonged effort that imaging a movement requires in comparison with movement execution was probably the reason for this stronger activation.

### Object observation

Several brain imaging studies have shown that naming tools, generating action words in response to tool observation, and imaging manipulating objects determine the activation of area 44 (see [45] for review). Recently, Chao and Martin [46\*\*] showed that even the mere observation of tools activates the cortex within the precentral sulcus, most likely in area 44 (−42, +6, +23). In addition, activation was also found in the left inferior parietal lobule, in an area corresponding to the intraparietal sulcus. Control conditions, in which pictures of animals, faces and houses were presented, showed that activation in both these regions was stronger with tools than with the other stimuli. These data show that, as in monkeys [4\*], observation of an object without other requirements activates a parietal–premotor circuit formed by area AIP and area 44 (the most likely human homologue of area F5).

In another very recent fMRI study [47\*], a parietal–premotor circuit largely overlapping that described by Chao and Martin [46\*\*] was found to be active during observation of bidimensional objects. The activation was particularly strong when the task required object-related attention.

### Action observation

A further condition that activates area 44 is action observation ([28\*\*,40,48], see also [49\*]). Iacoboni *et al.* [40] tested volunteers while they observed a finger movement, a static finger, or a cross (observation conditions), and when they were required to move their finger in response to these stimuli (observation/execution conditions). The results showed that, in the task in which participants had to move the finger in response to the observation of a similar movement (imitation), the activation of area 44 (−50, +12, +12) was stronger

than in all other conditions. These findings demonstrate that an action observation/execution matching system exists in humans, and is used for imitation. Similar results were obtained independently by Nishitani and Hari [41\*] using magnetoencephalography.

Activation of area 44 during the observation of hand actions could be interpreted as reflecting verbal mediation. If this interpretation is correct, area 44 should be active regardless of the type of observed action and the effector used. On the other hand, if activation of area 44 reflects a specific localization of hand actions, the activation should be absent when observing actions of other effectors, such as the foot. A recent fMRI experiment [28\*\*] showed that during the observation of hand actions there was an activation in area 44 (−64, +12, +20) and in ventral area 6 (see above). Foot action observation determined the activation of a dorsal site in area 6 (−32, −8, +64), but no activation of area 44. These results demonstrate that action observation activates somatotopically organized sectors of the premotor cortex, and that verbal mediation is not responsible for the activation of area 44 during hand and mouth action observation.

### Conclusions

The data reviewed here indicate that the vPM has both motor and cognitive functions. The motor functions transform the intrinsic properties of objects into hand actions and spatial locations into head and arm actions. The cognitive functions include space perception, action understanding and imitation.

There is also convincing evidence of a functional homology between monkey areas F5 and human area 44. Fewer data are available to establish the location of the possible human homologue of area F4. Preliminary evidence suggests that it is located in the ventral sector of area 6, approximately ventral to  $z=+50$ .

### References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Foerster, O: **The motor cortex in man in the light of Hughlings Jackson's doctrine.** *Brain* 1936, **59**:135-159.
  2. Von Bonin G, Bailey P: *The Neocortex of Macaca Mulatta.* Urbana IL: University of Illinois Press; 1947:136.
  3. Petrides M, Pandya DN: **Comparative architectonic analysis of the human and the macaque frontal cortex.** In *Handbook of Neuropsychology*, vol 9. Edited by Boller F, Grafman J. Amsterdam: Elsevier Science Ltd; 1994.
  4. Rizzolatti G, Luppino G: **The cortical motor system.** *Neuron* 2001, **31**:889-901.
 

An important review that summarizes the general organization of monkey cortical motor systems and discusses in detail some of the higher functions carried out by motor cortical areas. The authors suggest that the generation of internal representations of actions is central to cortical motor functions. External contingencies and motivational factors then determine whether these action representations are transformed into actual actions.
  5. Rizzolatti G, Berti A, Gallese V: **Spatial neglect: neurophysiological bases, cortical circuits, and theories.** In *Handbook of*

- Neuropsychology*, vol 1, edn 2. Edited by Boller F, Grafman J. Amsterdam: Elsevier Science Ltd; 2000.
6. Rizzolatti G, Scandolara C, Matelli M, Gentilucci M: **Afferent properties of periacuate neurons in macaque monkeys. II. Visual responses.** *Behav Brain Res* 1981, **2**:147-163.
  7. Gentilucci M, Fogassi L, Luppino G, Matelli M, Camarda R, Rizzolatti G: **Functional organization of inferior area 6 in the macaque monkey. I. Somatotopy and the control of proximal movements.** *Exp Brain Res* 1988, **71**:475-490.
  8. Graziano MSA, Gross CG: **Multiple pathways for processing visual space.** In *Attention and Performance XVI. Information Integration in Perception and Communication*. Edited by Inui T, McClelland JL. Cambridge, MA: MIT Press; 1996:181-207.
  9. Graziano MSA, Reiss LAJ, Gross CG: **A neuronal representation of the location of nearby sounds.** *Nature* 1999, **397**:428-430.
  10. Fogassi L, Gallese V, Fadiga L, Luppino G, Matelli M, Rizzolatti G: **Coding of peripersonal space in inferior premotor cortex (area F4).** *J Neurophysiol* 1996, **76**:141-157.
  11. Kakei S, Hoffmann DS, Strick PL: **Direction of action is represented in the ventral premotor cortex.** *Nat Neurosci* 2001, **4**:1020-1025.  
In this elegant study, recordings were obtained from vPM neurons in a monkey trained to perform wrist movements in different directions, with the wrist in three different postures. The task dissociated muscle activity, direction of joint movement and direction of movement in space. The results showed that nearly all neurons encode the direction of movement in space, independent of arm posture.
  12. Kakei S, Hoffmann DS, Strick PL: **Muscle and movement representations in the primary motor cortex.** *Science* 1999, **285**:2136-2139.
  13. Rizzolatti G, Matelli M, Pavesi G: **Deficits in attention and movement following the removal of postarcuate (area 6) and prearcuate (area 8) cortex in macaque monkeys.** *Brain* 1983, **106**:655-673.
  14. Halligan PW, Marshall JC: **Left neglect for near but not far space in man.** *Nature* 1991, **350**:498-500.
  15. Berti A, Frassinetti F: **When far becomes near: re-mapping of space by tool use.** *J Cogn Neurosci* 2000, **12**:415-420.  
This lesion study shows dissociation between far and near space in a patient with peripersonal neglect. The novel result is that when the patient used a tool for acting in far space, the neglect extended from near to far space. This provides a nice demonstration of a dynamic action-dependent mapping of space in humans.
  16. Iriki A, Tanaka M, Obayashi S, Iwamura Y: **Self-images in the video monitor coded by monkey intraparietal neurons.** *Neurosci Res* 2001, **40**:163-173.  
An extremely interesting study that extends the previous discovery by the same authors that tool use remaps the peripersonal visual RFs in the monkey. In this study, remapping also occurs when the monkey observes the image of its hand in a video monitor.
  17. Maravita A, Husain M, Clarke K, Driver J: **Reaching with a tool extends visual-tactile interactions into far space: evidence from cross-modal extinction.** *Neuropsychology* 2001, **39**:580-585.
  18. Farnè A, Ladavas E: **Dynamic size-change of hand peripersonal space following tool use.** *Neuroreport* 2000, **11**:1645-1649.
  19. Di Pellegrino G, Ladavas E, Farnè A: **Seeing where your hands are.** *Nature* 1997, **388**:730.
  20. Ladavas E, di Pellegrino G, Farnè A, Zeloni G: **Neuropsychological evidence of an integrated visuotactile representation of peripersonal space in humans.** *J Cogn Neurosci* 1998, **10**:581-589.
  21. Maravita A, Spence C, Clarke K, Husain M, Driver J: **Vision and touch through the looking glass in a case of crossmodal extinction.** *Neuroreport* 2000, **11**:3521-3526.  
The authors of this study show cross-modal extinction in the peripersonal space of a patient with a right hemisphere lesion. The important finding is that cross-modal extinction is present also when the patient sees his hand reflected in a mirror. This suggests that the vision of the hand via a mirror activates a representation of peripersonal space, and not of extrapersonal space suggested by the distant visual image in the mirror.
  22. Colebatch JG, Deiber MP, Passingham RE, Friston KJ, Frackowiak RS: **Regional cerebral blood flow during voluntary arm and hand movements in human subjects.** *J Neurophysiol* 1991, **65**:1392-1401.
  23. Matelli M, Rizzolatti G, Bettinardi V, Gilardi MC, Perani D, Rizzo G, Fazio F: **Activation of precentral and mesial motor areas during the execution of elementary proximal and distal arm movements: a PET study.** *Neuroreport* 1993, **4**:1295-1298.
  24. Toni I, Schluter ND, Josephs O, Friston K, Passingham RE: **Signal-, set- and movement-related activity in the human brain: an event-related fMRI study.** *Cereb Cortex* 1999, **9**:35-49.
  25. Desmurget M, Gréa H, Grethe JS, Prablanc C, Alexander GE, Grafton ST: **Functional anatomy of nonvisual feedback loops during reaching: a positron emission tomography study.** *J Neurosci* 2001, **21**:2919-2928.
  26. Bremner F, Schlack A, Jon Shah N, Zafiris O, Kubischik M, Hoffmann KP, Zilles K, Fink GR: **Polymodal motion processing in posterior parietal and premotor cortex: a human fMRI study strongly implies equivalencies between humans and monkeys.** *Neuron* 2001, **29**:287-296.  
An interesting fMRI study showing the localization of polymodal tactile, visual and auditory areas in the human brain. A circuit similar to that previously delineated in the monkey is described. The areas involved are the human homologues of monkey areas VIP, SII and F4.
  27. Galati G, Committeri G, Sanes JN, Pizzamiglio L: **Spatial coding of visual and somatic sensory information in body-centred coordinates.** *Eur J Neurosci* 2001, **14**:737-746.
  28. Buccino G, Binkofski F, Fink GR, Fadiga L, Fogassi L, Gallese V, Seitz RJ, Zilles K, Rizzolatti G, Freund H-J: **Action observation activates premotor and parietal areas in a somatotopic manner: an fMRI study.** *Eur J Neurosci* 2001, **13**:400-404.  
In this important fMRI study, the authors show that action observation elicits a somatotopic pattern of activation in the premotor cortex and the posterior parietal lobe. They demonstrate that the mirror system is not limited to area 44/F5, but encompasses a variety of parietal-premotor circuits. The observation of foot actions does not activate area 44, thus excluding an internal speech interpretation of action observation-related activations.
  29. Paus T: **Location and function of the human frontal eye-field: a selective review.** *Neuropsychology* 1996, **34**:475-483.
  30. Rizzolatti G, Camarda R, Fogassi L, Gentilucci M, Luppino G, Matelli M: **Functional organization of inferior area 6 in the macaque monkey. II. Area F5 and the control of distal movements.** *Exp Brain Res* 1988, **71**:491-507.
  31. Gallese V, Fadiga L, Fogassi L, Rizzolatti G: **Action recognition in the premotor cortex.** *Brain* 1996, **119**:593-609.
  32. Rizzolatti G, Fadiga L, Gallese V, Fogassi L: **Premotor cortex and the recognition of motor actions.** *Cogn Brain Res* 1996, **3**:131-141.
  33. Fogassi L, Gallese V, Buccino G, Craighero L, Fadiga L, Rizzolatti G: **Cortical mechanism for the visual guidance of hand grasping movements in the monkey: a reversible inactivation study.** *Brain* 2001, **124**:571-586.  
A reversible inactivation study of areas F5ab, F5c and F1 in the monkey. The inactivation of F5ab produced a selective deficit of visually guided hand shaping, preceding grasping and hand posture not appropriate for the size and shape of the object. The inactivation of F5c (the sector where mirror neurons are located) determined only a motor slowing, the hand shaping being preserved. The inactivation of the hand field of area F1 produced a severe paralysis of contralateral finger movements with hypotonia. The specificity of the deficits following F5ab lesion indicates that this F5 sector plays a crucial role in visuomotor transformation for object interaction.
  34. Umiltà MA, Kohler E, Gallese V, Fogassi L, Fadiga L, Keysers C, Rizzolatti G: **'I know what you are doing': a neurophysiological study.** *Neuron* 2001, **32**:91-101.  
The authors recorded from mirror neurons in area F5c during observation of grasping and holding actions. In one condition, the monkey saw the whole action, in another only its beginning, the crucial part of the action (hand-object interaction) being hidden by a screen. The majority of the recorded neurons responded also in the hidden condition. This study implies that the motor representation of an action performed by others is generated in the observer premotor cortex, even when a visual description of the action is lacking.
  35. Krams M, Rushworth MF, Deiber MP, Frackowiak RS, Passingham RE: **The preparation, execution and suppression of copied movements in the human brain.** *Exp Brain Res* 1998, **120**:386-398.
  36. Binkofski F, Buccino G, Posse S, Seitz RJ, Rizzolatti G, Freund H: **A fronto-parietal circuit for object manipulation in man: evidence from an fMRI study.** *Eur J Neurosci* 1999, **11**:3276-3286.

37. Ehrsson HH, Fagergren A, Jonsson T, Westling G, Johansson RS, Forsberg H: **Cortical activity in precision- versus power-grip tasks: an fMRI study.** *J Neurophysiol* 2000, **83**:528-536.
- In this study, the authors used fMRI to compare cortical activity during force production when participants used a precision grip and a power grip. The finding of a selective activation of area 44 during precision grip is extremely interesting.
38. Ehrsson HH, Fagergren A, Forsberg H: **Differential fronto-parietal activation depending on force used in a precision grip task: an fMRI study.** *J Neurophysiol* 2001, **85**:2613-2623.
39. Gerardin E, Sirigu A, Lehericy S, Poline J-B, Gaymard B, Marsault C, Agid Y, Le Bihan D: **Partially overlapping neural networks for real and imagined hand movements.** *Cereb Cortex* 2000, **10**:1093-1104.
- The authors of this fMRI study show that during hand action execution and imaging of the same action, an activation of a common network of areas occurs. This network includes area 44. Furthermore, during imaging, there is an activation of areas around the main network, including area 45, the presupplementary motor area, and predPM.
40. Iacoboni M, Woods RP, Brass M, Bekkering H, Mazziotta JC, Rizzolatti G: **Cortical mechanisms of human imitation.** *Science* 1999, **286**:2526-2528.
41. Nishitani N, Hari R: **Temporal dynamics of cortical representation for action.** *Proc Natl Acad Sci USA* 2000, **97**:913-918.
- The temporal dynamics of cortical activation during execution, observation and online imitations of right hand actions were studied in humans using neuro-magnetic recordings. During execution, left area 44 was activated first; this activation was followed by activation of left primary motor cortex and later, by the right primary motor cortex. During imitation and observation conditions, the sequence of activations was similar, but it started from the left occipital cortex. Activation was always strongest during action imitation. The authors conclude that the left area 44 is the 'orchestrator' of the human hand mirror system.
42. Decety J, Perani D, Jeannerod M, Bettinardi V, Tadini B, Woods R, Mazziotta JC, Fazio F: **Mapping motor representations with positron emission tomography.** *Nature* 1994, **371**:600-602.
43. Grafton ST, Arbib MA, Fadiga L, Rizzolatti G: **Localization of grasp representations in humans by PET. II. Observation compared with imagination.** *Exp Brain Res* 1996, **112**:103-111.
44. Parsons LM, Fox PT, Hunter Down J, Glass T, Hirsch TB, Martin CC, Jerabek PA, Lancaster JL: **Use of implicit motor imagery for visual shape discrimination as revealed by PET.** *Nature* 1995, **375**:54-58.
45. Martin A, Chao LL: **Semantic memory and the brain: structure and processes.** *Curr Opin Neurobiol* 2001, **11**:194-201.
46. Chao LL, Martin A: **Representation of manipulable man-made objects in the dorsal stream.** *Neuroimage* 2000, **12**:478-484.
- This important fMRI study reports two experiments in which pictures of tools, animals, faces and houses were presented to normal subjects. The tasks were either to observe or to silently name the stimuli. The results showed that the presentation of man-made manipulable objects activate a parieto-premotor circuit formed by the rostral vPM and by a region within the intraparietal sulcus. The control experiment with silent naming excluded that rostral vPM activation was due to verbalization.
47. Schubotz RI, von Cramon DY: **Functional organization of the lateral premotor cortex: fMRI reveals different regions activated by anticipation of object properties, location, and speed.** *Cogn Brain Res* 2001, **11**:97-112.
- In this interesting fMRI study, subjects were asked to attend to object-specific, spatial, or temporal properties of the same bidimensional moving objects. The results show that a parieto-frontal circuit comprising rostral vPM and a region in the anterior intraparietal sulcus, previously found to be active during prehension movements, was most intensely activated during object-related attention.
48. Grèzes J, Costes N, Decety J: **Top-down effect of strategy on the perception of human biological motion: a PET investigation.** *Cogn Neuropsychol* 1998, **15**:553-582.
49. Rizzolatti G, Fogassi L, Gallese V: **Neurophysiological mechanisms underlying the understanding and imitation of action.** *Nat Rev Neurosci* 2001, **2**:661-670.
- The most complete review on the mirror system in monkeys and humans. It includes an in depth discussion of the possible neural mechanisms that are at the basis of action understanding and imitation.
50. Talairach J, Tournoux P: *Co-planar Stereotaxic Atlas of the Human Brain.* Stuttgart: Georg Thieme Verlag; 1988.
51. Picard N, Strick PL: **Imaging the premotor areas.** *Curr Opin Neurobiol* 2001, **11**:663-672
- The authors of this interesting review summarize the most recent findings on mesial, dorsal, and ventral motor areas. Of great importance is the notion, formed on the basis of human and monkey experiments, that the dorsal premotor cortex is formed by a caudal (dPM, or F2) and rostral (predPM, or F7) sector. The rostral sector appears to have strong similarities with the prefrontal areas.